

GEF Report: Loan 952. Capturing a new phase of unrest at Santorini volcano, Greece.

David Pyle, University of Oxford.

Abstract.

Loan 952 supported the deployment of three cGPS receivers to augment geodetic studies of the first episode of active deformation and seismicity of the modern era at Santorini volcano, Greece. The instruments were deployed from July to October 2011, and were used primarily to complete a dense re-occupation of the twelve Greek Military Triangulation Points on Santorini. Processing of these data, and comparison with the earlier occupations of the same network (in 1955 and in the 1980's), provide the critical evidence that the episode of inflation at Santorini which took place from January 2011 – July 2012 was indeed the first major episode of deformation at this volcano since 1955; and most probably since the last eruption, in 1950. A report of our analysis of the geodetic data, combined with an analysis of the InSAR data for the period 2010-2012 has recently been published in Nature Geoscience.

Background.


In early 2011, during fieldwork by a PhD student, Michelle Parks, we became aware that an episode of unrest was beginning at the Greek island volcano of Santorini. Santorini is a classical caldera volcano, which experienced a major explosive eruption about 3600 years ago. Since that time, activity at the volcano has slowly built a steep-sided edifice of dacite lava in the centre of the caldera. This edifice first broke the surface in an eruption about 2000 years ago, and over the past 500 years the volcanic island of Nea Kameni has enlarged significantly in the course of five major eruptions. Although the eruptions themselves have been well described (e.g. Fouque, 1879; Pyle and Elliott, 2006), very little has been recorded relating to the unrest of the volcano in between eruptions: in fact, most of the sparse information in the literature relates only to 'precursor' seismicity and rapid uplift that occurred within a few weeks to months of an eruption starting. Thus, from the outset it was apparent that the new phase of unrest was a great opportunity to understand what happens to large caldera volcanoes in between eruptions. Once we had been able to confirm that the seismicity within the caldera was also accompanied by significant ground deformation (as observed both by InSAR, and from the evidence of the small number of operating cGPS stations on Santorini in early 2011), we applied to NERC for Urgency Funding to support an intensive campaign of field measurements, and to the GEF for some cGPS receivers to augment our fieldwork.

The field-intensive phase of measurement ran from June to October 2011, and included both campaign-style GPS measurements, as well the installation (with Greek colleagues from the National Technical University of Athens, NTUA) of several new permanent cGPS receivers, and field measurements of soil-gas flux. The activity included both junior and senior researchers from Oxford; researchers from Greece, and, for a period in September 2011, an intensive phase of campaign measurements by our 4th-year undergraduate class, who were visiting Santorini as a part of their field training.

The primary objectives of the field campaign were to capture the early stages of a continuing phase of unrest on Santorini, Greece. NERC Urgency funding and the GEF loan enabled us to perform the following tasks:

- to re-occupy the Greek Military Triangulation Points on Santorini.
- to extend the network of continuous GPS receivers on Santorini and
- to complete field surveys of gas emission from New Kameni.

Figure 1 (below) shows the locations of the remaining Greek Military Triangulation Points on Santorini that we re-occupied during the field campaign.



Survey Procedure.

For an initial survey in July 2011 we reoccupied campaign sites previously established by Dr Andy Newman (Georgia Tech) and his colleagues from the University of Patras. These sites are described and located in a recent paper in *Geophysical Research Letters* by Newman et al (2012). These campaign sites involved a mixture of pillar and tripod measurements. In September 2011, we reoccupied 12 Hellenic Military pillars (Fig. 1). We used a pool of equipment to do this, with instruments from both GEF and NTUA. During this campaign, several sites were occupied using the GEF receivers and choke ring antennae, but we used pillar plates from NTUA to mount the antennas on the pillars rather than using the tripods as these were too short. Sites were occupied for a minimum of 3 hrs during the September campaign and 24 hrs during the July campaign. We recognise that an ideal campaign survey may require longer occupation periods but in the case of the September campaign we were confident that 3 hours was sufficient because we used GPS observations from 3 closely located permanent stations on Santorini (SNTR, WNRV and DSLN) in the processing and we have short baselines from these cGPS sites to the pillars. SNTR was set up by University of Oxford/NTUA in 2005; WNRV and DSLN were set up in July 2011. The September survey campaign was carried out with the assistance of the 4th year undergraduate class from Oxford. This was an excellent educational experience for the students, who were enthused by the opportunity to participate in a field campaign.

The equipment provided by GEF was in very good condition and the setup proved to be very easy, once we had been trained. Probably the most difficult (and most important) task was programming the Leica receivers. Fortunately Colin Kay took us through the parameter file setup and the setup of equipment in the field which was extremely helpful. In the field, the most time consuming task when using a tripod setup was the levelling, but this became easier with experience. Overall, we were very pleased with the instruments.

Data quality, Processing, Modelling and Interpretation.

Full details are presented in our recent paper (Parks et al., 2012), and we summarise the salient points below. We used chain-stacked InSAR scenes from Envisat and TerraSAR-X to track the large-scale deformation field of Santorini from January 2011- April 2012. We used a matlab script to invert the data for the best-fitting 'Mogi' source (spherical pressure source), and find that the best-fit solution to both InSAR and cGPS data (from fixed stations) is for a Mogi source at about 4.5 km depth, located north of Nea Kameni, which is inflating at a rate of $\sim 10^7 \text{ m}^3 \text{ yr}^{-1}$ (Fig. 2, below). The location of this Mogi source has not changed significantly since May 2011. This is consistent with the unrest being due to the intrusion of a pulse of magma beneath the caldera.




Fig. 2. Best fit solutions to the location of a spherical Mogi pressure source (a), and its depth and volume (b), derived from modelling of the deformation field of Santorini measured by InSAR. The best fit solution puts the source north of the Kameni islands (the volcano in the centre of the caldera), and at 4 – 6 km depth. (See Parks et al., 2012 for further details).

We used the re-occupation of the triangulation points to investigate whether, or not, this current phase of deformation is the first such event since the last eruption in 1950. Triangulation observations were carried out on Santorini by the Hellenic Military Geographic Service in 1955. Each angle was the subject of 12 independent measurements, and processed in 1955 to yield a solution for coordinates (lat, long, orthometric height). For comparison with the GPS data, we subtracted the geoidal height from the orthometric height. We occupied the twelve surviving monuments from 23-26 September 2011, and the data from these sites and from four continuous GPS sites on Santorini were analysed using Bernese v5.0 GPS software. Coordinates were estimated in an ambiguity-fixed solution, using absolute antenna calibration corrections and IGS final products. The datum was realised by means of minimum constraints imposed on IGS stations BORI, BUCU, GLSV, ISTA, MATE, NICO, NOT1, PENC, WTZR. The processing was compliant with current Center for Orbit Determination in Europe standards.

Due to the imprecision of distance measurements in 1955, the initial network contains an undetermined scale error. We account for this by taking the three points most distant from the centre of inflation, and subtracted from their 2011 coordinates the displacements calculated from our InSAR-constrained model for inflation. This is a small adjustment. We then found the best fit solution of the isotropic horizontal dilatation to the 1955 coordinates to the lengths of baselines in the adjusted 2011 coordinates. We adopt a conservative estimate of 20 mm for the uncertainty in positions of the original coordinates, based on past experience of comparison between triangulation surveys and GPS observations.

The results, which are shown in Fig. 3, below confirm that the amount of deformation measured across the triangulation network (from 1955-2011) is essentially completely accounted for by the deformation that has happened during the current episode of unrest. This is evidence for the pulsatory nature of magmatic intrusion events beneath long-lived and dormant volcanoes.




Fig. 3. Horizontal surface displacements of Santorini between 1955 and 2011 from triangulation and GPS data. (a) The yellow vectors show the expected displacements if the only deformation in this time interval is that due to the inflation detected since the beginning of 2011. The red vectors show the displacements of monuments from 1955-2011, calculated from measurements of angle made in 1955 and GPS coordinates measured in 2011. The white circles show the locations of continuous GPS sites used in GPS processing. The scale of the network in 1955 is determined using the points shown with black circles. (b) Blue vectors show the differences between the model and observed; these represent displacements of the monuments from 1955 until the end of 2010. Uncertainties are shown by the white circles. There is no evidence for any large-scale deformation from 1955-2010.

Conclusions and recommendations

Use of GEF loan equipment was instrumental in our capturing of a short pulse of magma intrusion beneath Santorini volcano, and the confirmation that pulses of this scale happen only rarely in between eruptions. We are indebted to the staff of the facility for their expediting of our proposal, the rapid shipping of the equipment, and their dedication in being available at short notice to offer training in the field.

Publications

Peer-reviewed papers:

Parks, MM, Biggs, J, England, P, Mather, TA, Nomikou, P, Palamartchouk, K, Papanikolaou, X, Paradissis, D, Parsons, B, Pyle, DM, Raptakis, C, Zacharis, V, (2012) 'Evolution of Santorini Volcano dominated by episodic and rapid fluxes of melt from depth', *Nature Geoscience*, 5, 749-754, doi: 10.1038/NGEO1562

DPhil Thesis:

Parks, M.M., 2013, Volcanic processes during eruption and unrest: combining satellite and ground-based monitoring at Galeras and Santorini volcanoes, Unpublished D.Phil. thesis, University of Oxford. Link to Oxford Research Archive: <http://bit.ly/OeooNh>

Press coverage (mentioning the GEF loan)

NERC Planet Earth Online (Feature article, Autumn 2012),
<http://planetearth.nerc.ac.uk/features/story.aspx?id=1273>

Press Releases (from Oxford University, Bristol and ESA) were widely picked up by online media and news agencies, and coverage included print (Metro, UK, September 10th), television (Al Jazeera news, September 9th) and radio (BBC World Service, September 10th). Mainstream online coverage included National Geographic (<http://news.nationalgeographic.com/news/2012/09/120912-magma-balloon-lava-santorini-volcano-science/>) and CBBC Newsround (<http://www.bbc.co.uk/newsround/19549050>).

Conference presentations on Santorini unrest (chronological).

Michelle Parks et al., Poster, Fall AGU, December 2011

Michelle Parks et al., Volcanic and Magmatic Studies Winter Meeting, January 2012 (Durham UK).

Michelle Parks et al., Invited talk, EPOS, University of Athens, Greece (January 2012).

David Pyle et al., Invited Talk, MeMoVolc meeting of the Special Scientific Committee, Santorini, Greece (March 2012).

Juliet Biggs et al., ESA meeting on Satellite Earth Observation for Geohazard Risk Management, Santorini, Greece (May 2012).

David Pyle et al., Science Highlights talk, NCEO Annual Conference, Nottingham (September 2012).

Michelle Parks et al., Talk, Wegener 2012 Conference, Strasbourg (September 2012).

Table 1. Locations of GPS stations occupied during the field campaign.

In September 2011, we reoccupied 12 Hellenic Military pillars, using a pool of equipment from both GEF and NTUA. Sites were occupied for a minimum of 3 hrs, and we also used GPS observations from 3 permanent stations (SNTR, WNRV and DSLN) in processing and we have short baselines from these cGPS sites to the pillars.

Station	Latitude (N)	Longitude (E)	Elevation (m)
S010	36.21252436,	25.21330701,	169.724
S021	36.24149660,	25.25506509,	274.442
S022	36.24160327,	25.23415153,	161.797
S023	36.24260140,	25.28445544,	88.054
S028	36.25357382,	25.20482795,	330.438
S029	36.26017076,	25.26526602,	112.637
S030	36.25577835,	25.25197147,	402.967
S031	36.26409242,	25.26368758,	95.186
S034	36.27102646,	25.25253956,	352.469
S035	36.27422935,	25.23009615,	176.528
S302	36.22069714,	25.27496485,	602.479
S304	36.27483270,	25.24207002,	366.029

Table 1

Equipment provided by GEF for Santorini 2011 campaign GPS surveys	
3	Choke ring antennas and connector cables
3	Leica receivers
3	Tripods
3	CF cards for receivers
1	Copy of Leica GeoOffice Software and Dongle
3	Solar Panels
3	Solar Regulators and battery cables
2	Alloy containers
2	Battery chargers
3	External batteries and cables